

Decoquate and Monensin

Type B Medicated Cattle Feed

Do Not Feed Undiluted

INDICATIONS

For the prevention of coccidiosis caused by *Eimeria bovis* and *E. zuernii*, and improved feed efficiency in cattle being fed in confinement for slaughter.

ACTIVE DRUG INGREDIENTS¹

Decoquate ^a	535.8 to 5,440 g/ton
Monensin, USP ^b	500 to 80,000 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than	_____	%
Crude Fat, not less than	_____	%
Crude Fiber, not more than	_____	%
Calcium, not less than	_____	%
Calcium, not more than	_____	%
Phosphorus, not less than	_____	%
Potassium, not less than	_____	%
Salt ² , not less than	_____	%
Salt ² , not more than	_____	%
Sodium ³ , not less than	_____	%
Sodium ³ , not more than	_____	%

²If added.

³Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

MIXING DIRECTIONS

Mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of complete Type C medicated cattle feed containing 12.9 to 90.8 grams decoquate and 5 to 30 grams monensin (to provide 22.7 mg decoquate per 100 lb of body weight/day, 50 to 360 mg monensin/head/day). The Type C medicated feed is to be fed for the prevention of coccidiosis and improved feed efficiency during periods of exposure to coccidiosis or when it is likely to be a hazard. Feed only to cattle being fed in confinement for slaughter. Feed continuously as the sole ration. The following table provides examples of mixing and feeding rates.

Type B, grams/ton		Animal Wt. (lbs)	Lbs. Feed Per Head Per Day	Lbs. Type B per Ton of Feed	Lbs. Non-medicated Feed per Ton	Type C, grams/ton	
Decoquinatate	Monensin					Decoquinatate	Monensin
1136	1000	500	10	40	1960	22.7	20
2271	2015	500	15	13.3	1986.7	15.1	13.4
1136	715	700	14	40	1960	22.7	14.3
2271	1430	700	21	13.3	1986.7	15.1	9.5

CAUTION

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin horses has been fatal. Monensin medicated cattle feed is safe only for use in cattle. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of monensin may be fatal to cattle. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. Bentonite should not be used in decoquinatate feeds.

WARNINGS

Do not feed to animals producing milk for food. Do not feed to lactating dairy cattle.

When mixing and handling monensin use protective clothing, impervious gloves and a dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water.

Approved by FDA under ANADA # 200-652

**MANUFACTURED BY
BLUE BIRD FEED MILL
Any town, USA 12345**

Net Weight lb (kg) on bag or bulk

Expiration Date: 30 days after date of manufacture

Lot No. _____

¹Actual label must bear the single concentration of each drug.

^aSourced from Deccox[®], NADA # 039-417

^bSourced from Monovet[®], ANADA # 200-639

Deccox is a registered trademark of Zoetis Inc.

Monovet is a registered trademark of Huvepharma EOOD.