

Decoquinat e, Monensin, and Tylosin Type C Medicated Complete Feed

(decoquinat e and monensin and tylosin Type C medicat ed feed)

Caution: Federal law restricts medicat ed feed containing this veterinar y feed directive (VFD) drug to use by or on the order of a licens ed veterinar ian.

INDICATIONS

For the prevention of coccidiosis caused by Eimeria bovis and E. zuernii, for improved feed efficiency, and for reduction of incidence of liver abscesses associated with Fusobacterium necrophorum and Arcanobacterium pyogenes in cattle fed in confinement for slaughter.

ACTIVE DRUG INGREDIENTS¹

| | |
|---|---------------------------------|
| Decoquinat e ^a | 13.6 to 27.2 g/ton ² |
| Monensin ^b | 5 to 30 g/ton ² |
| Tylosin (as tylosin phosphate) ^c | 8 to 10 g/ton ² |

GUARANTEED ANALYSIS

| | |
|--|---------|
| Crude Protein, not less than..... | _____ % |
| Crude Fat, not less than..... | _____ % |
| Crude Fiber, not more than..... | _____ % |
| Calcium, not less than..... | _____ % |
| Calcium, not more than..... | _____ % |
| Phosphorus, not less than..... | _____ % |
| Potassium, not less than..... | _____ % |
| Salt ³ , not less than..... | _____ % |
| Salt ³ , not more than..... | _____ % |
| Sodium ⁴ , not less than..... | _____ % |
| Sodium ⁴ , not more than..... | _____ % |

³If added.

⁴Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed only to cattle being fed in confinement for slaughter. Feed continuously as the sole ration to provide 22.7 mg of decoquinat e per 100 lb. of body weight per day, 50 to 360 mg of monensin per head per day, and 60 to 90 mg tylosin per head per day as monensin sodium and as tylosin phosphate. Feed for at least 28 days during periods of exposure to coccidiosis or when it is likely to be a hazard. The following table provides examples of feeding rates and amount of drug per ton of feed:

| Animal Wt. (lbs.) | Feed Intake (% Body Wt) | Lbs. Feed Per Head /Day | Decoquinatate (grams/ton) | Monensin, 100 mg/head/day (grams/ton) | Tylosin | |
|----------------------|----------------------------|----------------------------|------------------------------|---|-------------|-----------|
| | | | | | mg/head/day | grams/ton |
| 500 | 2.8 | 14.0 | 16.2 | 14.3 | 60 | 8.6 |
| 700 | 2.5 | 17.5 | 18.2 | 11.4 | 70 | 8.0 |
| 900 | 2.2 | 19.8 | 20.6 | 10.1 | 90 | 9.1 |

CAUTION

Do not allow horses or other equines access to formulations containing monensin. Ingestion of monensin by equines has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result.

If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Bentonite should not be used in decoquinatate feeds.

WARNINGS

Warning: Do not feed to animals producing milk for human food. Do not feed to lactating dairy cattle. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

When mixing and handling monensin use protective clothing, impervious gloves and a dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water.

Approved by FDA under ANADA # 200-653

MANUFACTURED BY
BLUE BIRD FEED MILL
 Any town, USA 12345

Net Weight lb (kg) on bag or bulk

Lot No. (if applicable) _____

¹Final printed label on formulated Type C medicated feed must bear a single concentration for each drug.

²Concentrations (grams per ton) must be such that feed delivers a total intake of 22.7 mg decoquinatate per 100 pounds body weight per day, 50 to 360 mg monensin per head per day, and 60 to 90 mg tylosin per head per day. Indicate in feeding directions amount of medicated feed to be consumed daily.

^aSourced from Deccox[®], NADA # 039-417

^bSourced from Monovet[®], ANADA # 200-639

^cSourced from Tylovet[®], ANADA # 200-484

Deccox is a registered trademark of Zoetis Inc.

Monovet and Tylovet are registered trademarks of Huvepharma EOOD.