

# Bio-Cox<sup>®</sup>

---

## Granular 60

**Brand of Salinomycin Sodium  
Type A Medicated Article**

**Net wt 50 LB (22.68 kg)**



# Bio-Cox<sup>®</sup>

## Granular 60

### Brand of Salinomycin Sodium Type A Medicated Article

For use in broiler, roaster, replacement  
(breeder and layer) chickens, and quail only

**Active drug ingredient** Salinomycin equivalent to 60 grams salinomycin sodium activity per pound

**For use in manufacturing medicated feeds only**

**Broiler,  
roaster and  
replacement  
(breeder  
and layer)  
chickens**

Use Bio-Cox 60 Granular to provide salinomycin sodium concentrations ranging from 40-60 g/ton (0.0044% to 0.0066%) for the prevention of coccidiosis in broiler, roaster and replacement chickens caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti* and *E. mivati*. The dosage should be adjusted to meet the severity of the coccidial challenge, which varies with environmental and management conditions.

**Quail**

Use Bio-Cox 60 Granular to provide salinomycin sodium concentrations of 50 g/ton (0.0055%) for the prevention of coccidiosis in quail caused by *Eimeria dispersa* and *E. lettyae*.

**Mixing  
directions**

Thoroughly mix Bio-Cox 60 Granular with non-medicated feed to provide the level of salinomycin sodium per ton as indicated in the table below.

Salinomycin Sodium Activity (grams per ton)	Amount of Bio-Cox 60 Granular Type A Medicated Article (60 g/lb) Per Ton of Feed	
	lb	grams
40	0.67	304
45	0.75	341
50	0.83	377
55	0.92	418
60	1.00	454

**Feeding  
directions**

Feed continuously as the sole ration

**Important**

Must be thoroughly mixed in feeds before use. Not approved for use with pellet binders.

**Caution**

May be fatal if accidentally fed to adult turkeys or to horses.

**Warning**

**Do not feed to laying hens producing eggs for human consumption. When mixing and handling Bio-Cox 60 Granular premix, use protective clothing, impervious gloves, eye protection, and an approved dust respirator. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water.**

Store below 25°C (77°F), excursions permitted to 37°C (99°F)  
Restricted drug (California) - Use only as directed. Not for human use.  
**NADA #128-686, Approved by FDA**



Distributed by:  
Huvepharma, Inc.  
Peachtree City, GA 30269

Made in USA

Take Time



Observe Label  
Directions

HUVEPHARMA and Bio-Cox  
are registered trademarks of  
Huvepharma EOOD.

Net wt 50 LB (22.68 kg)

P08-400310  
Rev. 06-2016