

12 3/4"

6 3/4"

16"

6 3/4"

3 5/8"

CORID®

(amprolium)

Type A Medicated Article 25%

INDICATIONS: An aid in the prevention and treatment of coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves.

WARNING

Keep this and all drugs out of the reach of children. NOT FOR HUMAN USE.

The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report adverse effects, obtain an SDS or for assistance, contact 1-877-994-4883

Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

CAUTION: For satisfactory diagnosis, a microscopic examination of the feces should be done by a veterinarian or diagnostic laboratory before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

Do not use in feeds containing bentonite.

Use as the sole source of amprolium.

Restricted Drug – Use only as directed (California).

MIXING DIRECTIONS: CORID 25% should be thoroughly and evenly mixed in the feed. CORID 25% may be used to manufacture a Type B medicated feed with a concentration of 2.5% amprolium or a Type C medicated feed in the concentration range of 0.0125% to 1.25% amprolium.

- Use 200 pounds of CORID 25% per ton of feed to produce Type B medicated feed containing 2.5% amprolium.
- Use 1 pound of CORID 25% per ton of feed to produce Type C medicated feed containing 0.0125% amprolium.*
- Use 100 pounds of CORID 25% per ton of feed to produce Type C medicated feed containing 1.25% amprolium.
- Use 10 pounds of 2.5% Type B per ton of feed to produce Type C medicated feed containing 0.0125% amprolium.
- Use 1000 pounds of 2.5% Type B per ton of feed to produce Type C medicated feed containing 1.25% amprolium.

AMOUNT AND DIRECTIONS FOR USE:

Prevention: 227 mg amprolium/100 lb (5 mg/kg) body weight per day for 21 days during periods of exposure of when experience indicates that coccidiosis is likely to be a hazard.

Treatment: 454 mg amprolium/100 lb (10 mg/kg) body weight per day for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed and all calves in the group should be treated.

* To aid in even distribution of CORID 25% in the finished feed, prepare a mixture of CORID 25% in a portion of complete feed ingredient before mixing into the finished ration. Blend the mixture with the remainder of the finished feed and mix thoroughly.

The proper amount of amprolium supplement is mixed with the amount of ration usually consumed in one day. The tables below give suggested examples for using Type C medicated feed.


21 DAY PREVENTIVE PROGRAM:

Body Weight (lb)	DMI (as % Body Weight)	Amount of Feed per Day in Pounds		
		0.0125% Amprolium	0.0167% Amprolium	0.025% Amprolium
100	227	4	3	2
200	454	8	6	4
300	681	12	9	6
400	908	16	12	8
500	1134	20	15	10
600	1361	24	18	12

5 DAY TREATMENT PROGRAM:

Body Weight (lb)	DMI (as % Body Weight)	Amount of Feed per Day in Pounds		
		0.025% Amprolium	0.033% Amprolium	0.05% Amprolium
100	454	4	3	2
200	908	8	6	4
300	1362	12	9	6
400	1816	16	12	8
500	2268	20	15	10
600	2722	24	18	12

TAKE TIME



OBSERVE LABEL DIRECTIONS

www.corid.com

CORID®
(amprolium)
Type A Medicated Article 25%

CORID®

(amprolium)

Type A Medicated Article 25% Coccidiostat

To be used in the preparation of medicated feed to aid in the prevention and treatment of coccidiosis in calves.

ACTIVE DRUG INGREDIENT: Amprolium 25%

INGREDIENTS: Corn Gluten Feed and Soybean Oil

SEE DIRECTIONS ON BACK PANEL

STORAGE: Store at or below 77°F (25°C), excursions permitted to 104°F (40°C). Avoid prolonged exposure to high humidity.

NADA 12-350
Approved by the FDA

Product 00032

www.corid.com

® CORID is a registered trademark of Huvepharma EOOD. All rights reserved.

Net Wt.: 50 lb (22.68 kg)

Manufactured By:
Huvepharma, Inc.
525 Westpark Dr. Ste 230
Peachtree City, GA 30265, U.S.A.



CORID®

(amprolium)

Type A Medicated Article 25%

LOT NO & EXP DATE

30" (Less 5/8" Stretch)

5 3/4"