

# CORID<sup>®</sup>



(amprolium)

## 1.25% Pellets

Type C Medicated Feed  
Coccidiostat for Calves

An aid in the prevention and treatment of coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves.

**ACTIVE DRUG INGREDIENT:** See Back Panel of Bag

**DIRECTIONS:** See Back Panel of Bag

**INGREDIENTS:** See Back Panel of Bag

**GUARANTEED ANALYSIS:** See Back Panel of Bag

**MANUFACTURED UNDER LICENSE BY:** See Back Panel of Bag



Item # 200

**Net Wt.: 10 lb (4.53 kg)**



1022-2223-01  
Rev. 11-2006

® CORID and the CATTLE HEAD logo are registered trademarks of Merial Limited.



# CORID<sup>®</sup>



(amprolium)

## 1.25% Pellets Type C Medicated Feed Coccidiostat for Calves

An aid in the prevention and treatment of coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves.

**Indications for Use:** An aid in the prevention and treatment of coccidiosis caused by *Eimeria bovis* and *E. Zurnii* in calves

**Active Ingredient:** Amprolium 5,675 mg/lb.

#### Guaranteed Analysis

Protein	not less than	12.0%
Fat	not less than	1.5%
Fiber	not more than	22.0%

#### Ingredients

Dehydrated Alfalfa Meal, Wheat Midds, Calcium Carbonate, Soybean Oil, Lignin Sulfate, Artificial and or Natural Flavor.

**Feeding Directions:** Feed this feed as the sole source of amprolium to calves. Feed the proper amount of Corid 1.25% with the ration that will be consumed each day.

**21 Day Prevention:** Feed in complete feed or top dress at the rate of 227 mg amprolium per 100 lb body weight (5 mg/kg) daily for 21 days. Feed in at least one pound of feed per head per day.

Daily Corid 1.25% Feed Consumption in Pounds with Varying Bodyweight

Body Weight	Amprolium per Day (mg)	Amount of Corid 1.25% per Day in Pounds
100	227	0.04
200	454	0.08
300	681	0.12
400	908	0.16
500	1134	0.20
600	1361	0.24

For a group of animals, use the total body weight of the group to arrive at the proper amount of feed required.

**5 Day Treatment:** Feed in complete feed or top dress at the rate of 454 mg amprolium per 100 lb body weight (10 mg/kg) daily for 5 days. Feed in at least one pound of feed per head per day. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely the rest of the group has been exposed, and all calves in the group should be treated.

Daily Corid 1.25% Feed Consumption in Pounds with Varying Bodyweight

Body Weight	Amprolium per Day (mg)	Amount of Corid 1.25% per Day in Pounds
100	454	0.08
200	908	0.16
300	1362	0.24
400	1816	0.32
500	2268	0.40
600	2722	0.48

For a group of animals, use the total body weight of the group to arrive at the proper amount of feed required.

**Warning:** Withdraw 24 hours before slaughter. Use as the sole source of Amprolium  
A withdrawal period has not been established for this product in preruminating calves.  
Do not use in calves to be processed for veal.

**Caution:** Do not use Amprolium in feed containing bentonite.

Manufactured For:  
Agri Laboratories, Ltd  
St Joseph, MO 64503

Net Weight 10 lb (4.53 kg)

 **AgriLabs<sup>®</sup>**  
Making A Healthy Difference<sup>®</sup>

® CORID and the CATTLE HEAD logo are registered trademarks of Merial Limited.