

TYPE A MEDICATED ARTICLE

Sacox[®]60

BRAND OF SALINOMYCIN SODIUM

For Use in Broiler Chickens Only

For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti* and *E. mivati*.

Active Drug Ingredient

Salinomycin equivalent to 60 grams salinomycin sodium activity per pound.

Ingredients

Calcium Carbonate, Mineral Oil, Calcium Silicate, Silicon Dioxide and Microtracers.

Medicated Premix

Sixty (60) grams salinomycin sodium activity per pound.

Important

Must be Thoroughly Mixed in Feeds Before Use. Not approved for Use With Pellet Binders.

Restricted Drug (California) - Use Only as Directed

Net Weight 50 lbs. (22.68 kg)

Lot No. and Expiry Date printed on bag gusset.

STORAGE: Store at or below 25°C (77°F), excursions permitted to 40°C (104°F).
Avoid prolonged exposure to high humidity.

5135-05
Rev. 07-14

For Use in Broiler Chickens Only

Medicated Premix

Equivalent to 60 grams salinomycin sodium activity per pound. For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti* and *E. mivati*.

Dose

Sacox 60[®] can be used to provide concentrations of salinomycin ranging from 40 to 60 g/ton. The dosage should be adjusted to meet the severity of the coccidial challenge, which varies with environmental and management conditions.

Mixing Directions

Thoroughly mix the correct amount of the premix according to the directions below with an amount of non-medicated feed to provide the level of salinomycin per ton of feed.

| Salinomycin Sodium Activity (grams per ton) | Amount of Sacox [®] Premix Per Ton of Feed |
|--|--|
| 40 | 11 ounces |
| 45 | 12 ounces |
| 50 | 13.5 ounces |
| 55 | 14.5 ounces |
| 60 | 1 pound |

Important

Must Be Thoroughly Mixed in Feeds Before Use. Not for Use With Pellet Binders.

Feeding Directions

Feed continuously as the only ration.

Warning

No withdrawal required. Do not feed to laying chickens. May be fatal if accidentally fed to adult turkeys or to horses.

Warning

When mixing and handling Sacox 60[®] premix, use protective clothing, impervious gloves, eye protection and an approved dust respirator. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water.

Distributed by:

Huvepharma, Inc.
Peachtree City, GA 30269



ANADA # 200-075,
Approved by FDA