



Sacox® 60

(salinomycin sodium)

DESCRIPTION

Sacox® 60 brand of *salinomycin sodium* is a Type A Medicated Article for the prevention of coccidiosis in broilers, roaster and replacement (breeder and layer) chickens.

INDICATIONS

Use Sacox 60 to provide salinomycin sodium concentrations ranging from 40-60 g/ton (0.0044% to 0.0066%) for the prevention of coccidiosis in broiler, roaster and replacement chickens caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti* and *E. mivati*. The dosage should be adjusted to meet the severity of the coccidial challenge, which varies with environmental and management conditions

MIXING DIRECTIONS

Thoroughly mix Sacox 60 with non-medicated feed to provide the level of salinomycin sodium per ton as indicated in the table below.

Salinomycin Sodium Activity (grams per ton)	Amount of Sacox 60 Type A Medicated Article (60 g/lb) Per Ton of Feed	
	lb	grams
40	0.67	304
45	0.75	341
50	0.83	377
55	0.92	418
60	1.00	454

WARNING

Do not feed to laying hens producing eggs for human consumption. When mixing and handling Sacox 60 premix, use protective clothing, impervious gloves, eye protection and an approved dust respirator. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water.

WITHDRAWAL PERIOD

Sacox has no withdrawal requirement

CAUTION

May be fatal if accidentally fed to adult turkeys or to horses

STORAGE

Store below 25°C (77°F), excursions permitted to 37°C (99°F)



Take Time



Observe Label
Directions



SACOX is a registered
trademark of Huvepharma.

HVP-SACOX-PDS-08.17