## **RECOMMENDATIONS:**

PROHIBIT (levamisole hydrochloride) is a broadspectrum anthelmintic and is effective against the following nematode infections in cattle and sheep: STOMACH WORMS: (Haemonchus, Trichostrongylus, Ostertagia) INTESTINAL WORMS: (Trichostrongylus, Cooperia, Nematodirus, Bunostomum, Oesophagostomum) [Chabertia-Sheep only] LUNGWORMS: (Dictyocaulus)

RESIDUE WARNING: Do not administer to cattle within 48 hours of slaughter for food. Do not administer to sheep within 72 hours of slaughter for food. To prevent residues in milk, do not administer to dairy animals of breeding age.

MF# L-6232-05

Rev. 06-10

® Registered Trademark of Agri Laboratories, Ltd.

Manufactured for: Agri Laboratories, Ltd. St. Joseph, MO 64503

LOT NO .:

EXP. DATE:

NDC 57561-232-05

## **PROHIBIT®**

(levamisole hydrochloride) Soluble Drench Powder Anthelmintic

FOR ORAL USE IN CATTLE AND SHEEP

Each packet contains 46.8 grams of levamisole hydrochloride activity.

For Animal Use Only Keep Out of Reach of Children Restricted Drug (California) -Use Only as Directed ANADA 200-225, Approved by FDA

Net Contents: 30 x 52 Gram Pouches



**DOSAGE AND ADMINISTRATION:** Refer to packet for complete dosage and administration instructions.

NOTE: Careful weight estimates are essential for proper performance of this product. Prepare solutions as needed. However, excess solutions may be stored in clean closed containers up to 90 days without loss of anthelmintic activity. Cattle and sheep maintained under conditions of constant helminth exposure may require retreatment within two to four weeks after the first treatment.

**CAUTION:** Muzzle foam may be observed. However, this reaction will disappear within a few hours. If this condition persists, a veterinarian should be consulted. Follow recommended dosage carefully. Consult veterinarian before using in severely debilitated animals.

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

Store between 15°-30°C (59-86°F).

