

Tylovet® 10

(tylosin phosphate)



NET WEIGHT 50 LBS (22.68 KG.)

Approved by FDA under ANADA# 200-484

Manufactured by: Biovet Ad | 39 Petar Rakov Str. 4450 Peshtera, Bulgaria
Distributed by: Huvepharma, Inc.

For Use In Swine and Beef Cattle Feeds Only

Type A Medicated Article

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For Use in Swine and Beef Cattle Feeds Only

Do Not Feed Undiluted

Swine:

- For control of porcine proliferative enteropathies (PPE, ileitis) associated with *Lawsonia intracellularis*.
- For control of swine dysentery associated with *Brachyspira hyodysenteriae*.
- For the reduction in severity of effects of atrophic rhinitis.
- For treatment and control of swine dysentery associated with *Brachyspira hyodysenteriae* immediately after medicating with tylosin in drinking water.
- For control of porcine proliferative enteropathies (PPE, ileitis) associated with *Lawsonia intracellularis* immediately after medicating with tylosin in drinking water.

Beef Cattle:

- For reduction of incidence of liver abscesses associated with *Fusobacterium necrophorum* and *Arcanobacterium pyogenes*.

Storage

Store at room temperature, 25°C (77°F). Avoid moisture.

Not to be used after the date printed on the bottom of the bag.

To report adverse effects, access medical information, or obtain additional product information, call 1-877-426-7765. View product label <http://www.huvepharma.us/media/documents/products/labels/TYLOVET-100-Label.PDF>.

DIRECTIONS FOR USE (Read All Directions Carefully Before Mixing and Feeding):

Active Drug Ingredient: Tylosin (as tylosin phosphate) 10 grams per pound

Ingredients: Rice Hulls, Calcium Carbonate and Mineral Oil.

Crude Protein, Minimum 1.8%
Crude Fat, Minimum 1.9%
Crude Fiber, Maximum 35.5%
Calcium (Ca), Minimum 9.5%
Calcium (Ca), Maximum 11.4%
Phosphorous (P), Minimum 0.1%

This product has been formulated to contain at least 25 percent nutrient content.

Important: Must be thoroughly mixed in feeds before use.

CAUTION: Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

Swine

Mixing Directions for Medicated Feed for control

Usage	Tylovet 10 per Ton of Complete Feed	Tylosin per Ton of Complete Feed
For control of PPE (ileitis)	10 lbs --or-- 10 lbs then 4 lbs	100 g for 21 days --or-- 100 g for at least 21 days --then-- 40 g to market
For control of swine dysentery	10 lbs --then-- 4 lbs	100 g for at least 21 days --then-- 40 g to market
For reduction in severity of effects of atrophic rhinitis	10 lbs	100 g
For treatment and control of swine dysentery immediately after medicating with tylosin in drinking water	4 to 10 lbs	40 to 100 g (2 to 6 weeks)
For control of PPE (ileitis) immediately after medicating with tylosin in drinking water	4 to 10 lbs	40 to 100 g (2 to 6 weeks)

Beef Cattle

Mixing Directions for Medicated Feed

Usage	Tylovet 10 per Ton of Complete Feed	Tylosin per Ton of Complete Feed
For reduction of incidence of liver abscesses in beef cattle	0.8 to 1.0 lb	8 to 10 g

Huvepharma and Tylovet are registered trademarks of Huvepharma EOOD.

Tylovet® 40

(tylosin phosphate)



NET WEIGHT 50 LBS (22.68 KG.)

Approved by FDA under ANADA# 200-484

Manufactured by: Biovet Ad | 39 Petar Rakov Str. 4450 Peshtera, Bulgaria
Distributed by: Huvepharma, Inc.

For Use In Swine and Beef Cattle Feeds Only

Type A Medicated Article

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For Use in Swine and Beef Cattle Feeds Only

Do Not Feed Undiluted

Equivalent to 40 g Tylosin per Pound

Swine:

- For the reduction in severity of effects of atrophic rhinitis.
- For control of swine dysentery associated with *Brachyspira hyodysenteriae*.
- For the treatment and control of swine dysentery associated with *Brachyspira hyodysenteriae* immediately after medicating with **Tylovet Soluble** (tylosin tartrate) drinking water.
- For control of porcine proliferative enteropathies (PPE, ileitis) associated with *Lawsonia intracellularis*.
- For control of porcine proliferative enteropathies (PPE, ileitis) associated with *Lawsonia intracellularis* immediately after medicating with **Tylovet Soluble** (tylosin tartrate) drinking water.

Beef Cattle:

- For reduction of incidence of liver abscesses associated with *Fusobacterium necrophorum* and *Arcanobacterium pyogenes*.

Storage:

Store at room temperature, 25°C (77°F). Avoid moisture.

Not to be used after the date printed on the bottom of the bag.

To report adverse effects, access medical information, or obtain additional product information, call 1-877-426-7765. View product label <http://www.huvepharma.us/media/documents/products/labels/TYLOVET-100-Label.PDF>.

DIRECTIONS FOR USE (Read All Directions Carefully Before Mixing and Feeding):

Tylovet® 40 Type A Medicated Article

Active Drug Ingredient: Tylosin (as tylosin phosphate) 40 g per lb

Ingredients: Roughage products, pregelatinized starch and dipotassium phosphate.

Caution: To ensure adequate mixing, an intermediate blending step should be used prior to manufacturing a complete feed. Do not use in any liquid feed containing sodium metabisulfite or in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

Warning: Tylovet 40 may be irritating to unprotected skin and eyes. When mixing and handling Tylovet 40 use protective clothing, impervious gloves and a dust respirator. In case of accidental eye exposure, flush eyes with plenty of water. Exposed skin should be washed with plenty of soap and water. Remove and wash contaminated clothing. Seek medical attention if irritation becomes severe or persists. The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report adverse effects, access medical information, or obtain additional product information, call 1-877-426-7765.

For the reduction in severity of effects of atrophic rhinitis. Feed 100 g of tylosin per ton (2.5 pounds Tylovet 40 per ton) of complete feed. Feed continuously as the sole ration.

For control of swine dysentery. Feed 100 g of tylosin per ton (2.5 pounds Tylovet 40 per ton) of complete feed for at least three weeks. Follow with 40 g tylosin per ton (1 pound Tylovet 40 per ton) of complete feed until pigs reach market weight.

For the treatment and control of swine dysentery. Feed 40 to 100 grams of tylosin (1.0 to 2.5 pounds of Tylovet 40) per ton of complete feed for 2 to 6 weeks immediately after medicating with 250 mg tylosin (as Tylovet Soluble) per gallon in drinking water for 3 to 10 days .

For control of porcine proliferative enteropathies (PPE, ileitis). Feed 100 g of tylosin per ton (2.5 pounds Tylovet 40 per ton) of complete feed for 21 days. Alternatively, feed 100 g of tylosin per ton (2.5 pounds Tylovet 40 per ton) of complete feed for at least 3 weeks, followed by 40 g tylosin per ton of complete feed until pigs reach market weight. Alternatively, feed 40 to 100 grams of tylosin (1.0 to 2.5 pounds of Tylovet 40) per ton of complete feed for 2 to 6 weeks immediately after medicating with 250 mg tylosin (as Tylovet Soluble) per gallon in drinking water for 3 to 10 days. Feed continuously as the sole ration when feeding Tylovet.

NOTICE: Organisms vary in their degree of susceptibility to any chemotherapeutic. If no improvement is observed after recommended treatment, diagnosis and susceptibility should be reconfirmed.

Mixing and Feeding Directions for Beef Cattle Feeds For reduction of incidence of liver abscesses associated with <i>Fusobacterium necrophorum</i> and <i>Arcanobacterium pyogenes</i>	Tylovet 40 per Ton of Type C Feed	Tylosin Per Ton of Type C Feed
	0.2 to 0.25 lbs To be fed so that each animal receives not more than 90 mg per head per day and not less than 60 mg per head per day. Feed continuously as the sole ration.	8 to 10 g
Mixing Directions for Liquid Type B Cattle Feeds	LIMITATIONS: 1. pH must be between 4.5 and 6.0. 2. For liquid feeds stored in recirculating tank systems: Recirculate immediately prior to use for not less than 10 minutes, moving not less than 1 percent of the tank contents per minute from the bottom of the tank to the top. Recirculate daily as described even when not used. 3. For liquid feeds stored in mechanical, air, or other agitation - type tank systems: Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily as described even when not used. 4. Liquid Type B Cattle Feeds must bear an expiration date of 31 days after the date of manufacture.	

Huvepharma and Tylovet are registered trademarks of Huvepharma EOOD.

Tylovet® 100

(tylosin phosphate)



NET WEIGHT 50 LBS (22.68 KG.)

Approved by FDA under ANADA# 200-484

Manufactured by: Biovet Ad | 39 Petar Rakov Str. 4450 Peshtera, Bulgaria
Distributed by: Huvepharma, Inc.

For Use In Swine and Beef Cattle Feeds Only

Type A Medicated Article

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For Use in Swine and Beef Cattle Feeds Only

Do Not Feed Undiluted

Equivalent to 100 g Tylosin per Pound

Swine:

- For the reduction in severity of effects of atrophic rhinitis.
- For control of swine dysentery associated with *Brachyspira hyodysenteriae*.
- For the treatment and control of swine dysentery associated with *Brachyspira hyodysenteriae* immediately after medicating with **Tylovet Soluble** (tylosin) drinking water.
- For control of porcine proliferative enteropathies (PPE, ileitis) associated with *Lawsonia intracellularis*.
- For control of porcine proliferative enteropathies (PPE, ileitis) associated with *Lawsonia intracellularis* immediately after medicating with **Tylovet Soluble** (tylosin) drinking water

Beef Cattle:

- For reduction of incidence of liver abscesses associated with *Fusobacterium necrophorum* and *Arcanobacterium pyogenes*.

Storage:

Store at room temperature, 25°C (77°F). Avoid moisture.

Not to be used after the date printed on the bottom of the bag.

To report adverse effects, access medical information, or obtain additional product information, call 1-877-426-7765. View product label <http://www.huvepharma.us/media/documents/products/labels/TYLOVET-100-Label.PDF>.

DIRECTIONS FOR USE (Read All Directions Carefully Before Mixing and Feeding):

Type A Medicated Article

Active Drug Ingredient: Tylosin (as tylosin phosphate) 100 g per lb

Ingredients: Roughage products, pregelatinized starch and dipotassium phosphate.

Caution: To ensure adequate mixing, an intermediate blending step should be used prior to manufacturing a complete feed. Do not use in any liquid feed containing sodium metabisulfite or in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

Warning: **Tylovet 100** may be irritating to unprotected skin and eyes. When mixing and handling **Tylovet 100** use protective clothing, impervious gloves and a dust respirator. In case of accidental eye exposure, flush eyes with plenty of water. Exposed skin should be washed with plenty of soap and water. Remove and wash contaminated clothing. Seek medical attention if irritation becomes severe or persists. The material safety data sheet (MSDS) contains more detailed occupational safety information. To report adverse effects, access medical information, or obtain additional product information, call 1-877-426-7765.

For the reduction in severity of effects of atrophic rhinitis. Feed 100 g of tylosin per ton (1.0 pound **Tylovet 100** per ton) of complete feed. Feed continuously as the sole ration.

For control of swine dysentery. Feed 100 g of tylosin per ton (1.0 pound **Tylovet 100** per ton) of complete feed for at least three weeks. Follow with 40 g tylosin per ton (0.4 pound **Tylovet 100** per ton) of complete feed until pigs reach market weight.

For the treatment and control of swine dysentery. Feed 40 to 100 grams of tylosin (0.4 to 1.0 pounds of **Tylovet 100**) per ton of complete feed for 2 to 6 weeks immediately after medicating with 250 mg tylosin (as **Tylovet Soluble**) per gallon in drinking water for 3 to 10 days .

For control of porcine proliferative enteropathies (PPE, ileitis). Feed 100 g of tylosin per ton (1.0 pound **Tylovet 100** per ton) of complete feed for 21 days. Alternatively, feed 100 g of tylosin per ton (1.0 pound **Tylovet 100** per ton) of complete feed for at least three weeks, followed by 40 g tylosin per ton of complete feed until pigs reach market weight. Alternatively, feed 40 to 100 grams of tylosin (0.4 to 1.0 pounds of **Tylovet 100**) per ton of complete feed for 2 to 6 weeks immediately after medicating with 250 mg tylosin (as **Tylovet Soluble**) per gallon in drinking water for 3 to 10 days. Feed continuously as the sole ration when feeding Tylovet.

NOTICE: Organisms vary in their degree of susceptibility to any chemotherapeutic. If no improvement is observed after recommended treatment, diagnosis and susceptibility should be reconfirmed.

Mixing and Feeding Directions for Beef Cattle Feeds For reduction of incidence of liver abscesses associated with <i>Fusobacterium necrophorum</i> and <i>Arcanobacterium pyogenes</i>	Tylovet 100 per Ton of Type C Feed	Tylosin Per Ton of Type C Feed
	0.08 to 0.10 lbs To be fed so that each animal receives not more than 90 mg per head per day and not less than 60 mg per head per day. Feed continuously as the sole ration.	8 to 10 g
Mixing Directions for Liquid Type B Cattle Feeds	LIMITATIONS: 1. pH must be between 4.5 and 6.0. 2. For liquid feeds stored in recirculating tank systems: Recirculate immediately prior to use for not less than 10 minutes, moving not less than 1 percent of the tank contents per minute from the bottom of the tank to the top. Recirculate daily as described even when not used. 3. For liquid feeds stored in mechanical, air, or other agitation - type tank systems: Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily as described even when not used. 4. Liquid Type B Cattle Feeds must bear an expiration date of 31 days after the date of manufacture.	

Huvepharma and Tylovet are registered trademarks of Huvepharma EOOD.