## CORD (amprolium) 9.6% Oral Solution Coccidiostat

Net Contents: 1 gal. (128 fl. oz) 3,785 mL) Active Ingredient: amprolium ......9.6%

NADA 013-149, Approved by FDA

Distributed by: Huvepharma, Inc. Peachtree City, GA 30269



INDICATIONS: An aid in the treatment and prevention of coccidiosis caused by Eimeria bovis and E. zurnii in calves. For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

## DOSAGE AND ADMINISTRATION:

**5-Day Treatment Protocol** Daily Dosage: 10 mg amprolium/kg (10 mg per 2.2 lb body weight)

21-Dav Prevention Protocol Daily Dosage: 5 mg amprolium/kg (5 mg per 2.2 lb body weight)

**USE DIRECTIONS:** There are **96 mg** of amprolium in every **1 mL** of CORID 9.6% Solution.

1 fl oz = 29.57 mL 8 fl oz = ½ pint (236.56 mL) 1 fl oz = 2 measuring tablespoons 16 fl oz = 1 pint (473.12 mL)

## **IN DRINKING WATER**

Mix CORID daily with fresh drinking water.

**5-DAY TREATMENT:** Add CORID 9.6% Oral Solution to drinking water at the rate of 16 fl oz/100 gal. At the usual rate of water consumption this will provide an intake of approximately 10 mg amprolium/kg (2,2 lb) body weight. Offer this solution as the only source of water for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

**21-DAY PREVENTION:** During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add CORID 9.6% Oral Solution to drinking water at the rate of 8 fl oz/ 100 gal. At usual rates of water consumption this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the the only source of water for 21 days.

HUVEPHARMA	GRAPHIC	ARTWORK	APPROVAL
------------	---------	---------	----------

ITEM CODE: KCP-S00010	VERSIO	N	COLORS	NOTES	Marketing approval:	Regulat
PRODUCT: Corid 9.6% Gallon Bottle Label SPEC#: N/A	INITIALS: VERSION - DATE:	xx KCP-S00010 Draft 2 Rev: 06-2017	К Р 485	SITE/SUPPLIER: Huvepharma SOFTWARE: ADOBE	COMPLETE NAME + DATE + SIGNATURE:	COMPLETE N
COUNTRY: DOMESTIC (US)						
Dimensions: mm = inches = 4.875 x 17			<ul> <li>Please use the official PANTONE® (P) matching</li> <li>system for an accurate color representation.</li> </ul>			

Artwork Approved: Theresa Palmer 28-JUL-2017

## **AS A DRENCH**

CORID drench solutions may be stored in a clean, closed, labeled container for up to 3 days 5-DAY TREATMENT: Add 3 fl oz CORID 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2 lb) body weight. Give daily for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

**21-DAY PREVENTION:** During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1 ½ fl oz of CORID 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/ kg (2.2 lb) body weight. Give daily for 21 days.

**RESIDUE WARNINGS:** Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for yeal.

WARNING: NOT FOR HUMAN USE. Keep this and all drugs out of the reach of children. The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report adverse reactions in users, to obtain more information, or to obtain an SDS, contact Huvepharma at 1-877-426-7765.

**PRECAUTIONS: FOR ORAL USE IN ANIMALS ONLY. MAY CAUSE EYE IRRITATION.** For irritations, flush with plenty of water; get medical attention.

**STORAGE:** Store between 5°-25°C (41°-77°F) with brief excursions to 40°C.

Benzoic acid 0.1% added as preservative.

© CORID is a registered trademark of Huvepharma EOOD. Lot Number and Expiration Date can be found on shoulder of bottle



KCP-S00010 Version: 01 Rev: 06-2017

<b>ulatory</b> approval:	PRINTER NOTES!
LETE NAME + DATE + SIGNATURE:	