

8.9535" CUTOFF 2 AROUND 17.907" CYLINDER .067" PLATES

SLIT 1/8" OFF EDGE OF EYEMARK

12.875" WEB WIDTH

IN DRINKING WATER

Mix CORID daily with fresh drinking water. Stock solution for water proportioners may be stored in a clean, closed, labeled container for up to 3 days.

5-DAY TREATMENT: Add CORID 20% Soluble Powder to drinking water at the rate of 4 oz/50 gal. One 10 oz pack to 125 gallons of water. For use in automatic water proportioners that meter 1 fl oz of stock solution per gallon of drinking water, add 1 bag (10 oz) in 1 gallon of water to make stock solution. At the usual rate of water consumption, this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Treatment should be on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add CORID 20% Soluble Powder to drinking water at the rate of 4 oz/100 gal. One 10 oz pack to 250 gallons of water. For use in automatic water proportioners that meter 1 fl oz of stock solution per gallon of drinking water, add 1/2 bag (5 oz) in 1 gallon of water to make stock solution. At usual rates of water consumption, this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 21 days.

AS A DRENCH

CORID drench solutions may be stored in a clean, closed, labeled container for up to 3 days.

5-DAY TREATMENT: Add 3 oz CORID 20% Soluble Powder to 1 qt (32 fl oz) of water and, with a dose syringe, give 1 fl oz of this solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2 lb) body weight. Give daily for 5 days. Treatment should be on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1 1/2 oz of CORID 20% Soluble Powder to 1 qt (32 fl oz) of water and, with a dose syringe, give 1 fl oz of this solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21 days.

Lot No & Exp Date ▼

CORID[®]

(amprolium)

20% Soluble Powder Coccidiostat

An aid in the treatment and prevention of bovine coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves

Net Contents: 10 oz. (283.5 grams)

Active Ingredient: amprolium.....20%

Distributed by:
Huvepharma, Inc.
Peachtree City, GA 30269

For questions, please call 1-877-426-7765
NADA 033-165, Approved by FDA



CORID[®] (amprolium) 20% Soluble Powder - FOR ANIMAL USE ONLY

INDICATION: An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves.

CAUTION: For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

USE DIRECTIONS: There are **200 mg** of amprolium in every **1 gram** of CORID 20% Soluble Powder.

5-Day Treatment Protocol Daily Dosage: 10 mg amprolium/kg (10 mg per 2.2 lb body weight)	21-Day Prevention Protocol Daily Dosage: 5 mg amprolium/kg (5 mg per 2.2 lb body weight)
-------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------

1 oz = 3 1/2 measuring tablespoonsful

1 oz = 28.35 grams

RESIDUE WARNING: Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

NOT FOR HUMAN USE. Restricted Drug (California) - Use only as directed. Keep this and all drugs out of the reach of children.

Storage: Store between 5°-25°C (41°-77°F) with brief excursions to 40°C (104°F). Avoid high humidity storage conditions.

The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report adverse reactions in users, to obtain more information, or to obtain a SDS, contact Huvepharma at 1-877-426-7765.

©CORID is a trademark of Huvepharma EOOD.



KCP-S00015
Version: 01
Rev: 06-2017

HUVEPHARMA GRAPHIC ARTWORK APPROVAL

ITEM CODE: xxxx-xxxx PRODUCT: Corid 20% DISPLAY TRAY SPEC#: N/A COUNTRY: DOMESTIC (US) Dimensions: mm = inches = 36" x 47-1/8"	VERSION INITIALS: TP VERSION - DATE: KCP-S00015 Version 01 Rev. 06-2017	COLORS K P 485 Please use the official PANTONE® (P) matching system for an accurate color representation.	NOTES SITE/SUPPLIER: Huvepharma SOFTWARE: ADOBE	Marketing approval: COMPLETE NAME + DATE + SIGNATURE:	Regulatory approval: COMPLETE NAME + DATE + SIGNATURE:	PRINTER NOTES!
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------	-----------------------------------------------------------------	------------------------------------------------------------------	-----------------------