

Serial #

Exp.

MORAXELLA BOVIS BACTERIN

Single Dose Pinkeye Bacterin

10 Doses / 20 mL



I-SITE XP®



Dosage: Shake Well. Adminster a 2 mL dose subcutaneously to cattle 2 months of age or older.

Caution: Store at 2 to 8°C (35 to 46°F). DO NOT FREEZE. Shake well before and during use. Use entire contents when first opened. Do not mix with other products. In case of anaphylactic reaction, administer epinephrine or atropine. Do not vaccinate within 21 days before slaughter. For use in animals only. In case of human exposure, contact a physician. This product has not been tested in pregnant animals.

© Registered Trademark of Huvepharma, Inc.

Manufactured by:
Addison Biological Laboratory, Inc.
Fayette, Missouri 65248



Distributed by:
Huvepharma, Inc.
Peachtree City, GA 30269

VLN 355/PCN 2772.00

RV0021

Composition: This vaccine contains formalin inactivated isolates of *Moraxella bovis*.

Indications: This product has been shown to be effective for the vaccination of healthy cattle 2 months of age or older against pinkeye (infectious bovine keratoconjunctivitis). The duration of immunity against *Moraxella bovis* is at least 120 days when administered at 5 months of age.

For more information regarding efficacy and safety data, see productdata.aphis.usda.gov.

Technical Inquiries or Adverse Events:
800-331-2530

