

Monensin and Tylosin Medicated Cattle Feed – Feed Efficiency Type C Medicated Feed

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

INDICATIONS

For improved feed efficiency and reduction of incidence of liver abscesses caused by *Fusobacterium necrophorum* and *Arcanobacterium (Actinomyces) pyogenes* in cattle fed in confinement for slaughter.

ACTIVE DRUG INGREDIENTS

Monensin, USP ¹	5 to 40 g/ton**
Tylosin phosphate ²	8 to 10 g/ton**

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Non-Protein Nitrogen (NPN) ³ , not more than	_____	%
Crude Fat, not less than	_____	%
Crude Fiber, not more than	_____	%
Calcium, not less than	_____	%
Calcium, not more than	_____	%
Phosphorus, not less than	_____	%
Salt ⁴ , not less than.....	_____	%
Salt ⁴ , not more than	_____	%
Sodium ⁵ , not less than.....	_____	%
Sodium ⁵ , not more than	_____	%
Potassium, not less than	_____	%
Vitamin A ^{4,6} , not less than	_____	I.U./lb

³When added.

⁴If added.

⁵Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁶Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS


Feed continuously as sole ration to provide 50 to 480 mg monensin/hd/day and 60 to 90 mg/hd/day tylosin. No additional improvement in feed efficiency has been shown from feeding monensin at levels greater than 30 g/ton (360 mg monensin per head per day).

CAUTIONS


Do not allow horses or other equines access to feeds containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

WARNINGS



A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.



Approved by FDA under ANADA # 200-642

MANUFACTURED BY
BLUE BIRD FEED MILL
Anytown, USA 12345

Net Weight lb (kg) on bag or bulk

Lot Number: _____

**The medicated feed label must state a single drug concentration.

¹Sourced from Monovet[®], ANADA # 200-639

²Sourced from Tylovet[®], ANADA # 200-484

Monovet and Tylovet are registered trademarks of Huvepharma EOOD.