


NO COPY



Warning (Human Food)
Do not treat chickens or turkeys within 14 days of slaughter for food. Do not medicate chickens or turkeys producing eggs for human consumption.

Directions for Use:

	PROPORTIONER SOLUTION (1 oz/gal)	TANK MIX
.04% Solution	Add one pack to 1 gallon (3.8 liters)	Add one pack to 128 gallons
.025% Solution	Add one pack to 1.6 gallons (6.1 liters)	Add one pack to 206 gallons

MAKE FRESH SOLUTION DAILY. If improvement is not noted in 72 hours, consult your veterinarian. During treatment use only medicated water unless otherwise directed. For control of disease outbreaks medication should be initiated as soon as diagnosis is determined. Treated animals must actually consume enough medicated water to provide a therapeutic dose. Do not mix or administer in galvanized containers. Dispose of any waste or unused portions properly.

PRECAUTION: May cause toxic reactions unless drug is evenly mixed in water at dosages indicated and used according to label directions.

CORNER RADIUS

PoultrySulfa®

(sulfamerazine, sulfamethazine and sulfaquinoxaline)

Antimicrobial
For Oral Veterinary Use Only
CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

For Use in Drinking Water Only

As an aid in the control of coccidiosis and acute fowl cholera in chickens and acute fowl cholera and coccidiosis in turkeys, when caused by pathogens susceptible to sulfamerazine, sulfamethazine and sulfaquinoxaline.



Manufactured for:
Huvepharma, Inc.
525 Westpark Drive, Suite 230
Peachtree City, GA 30269

SOLUBLE POWDER

For Use in Chickens and Turkeys

THIS PACKET CONTAINS:
78 grams Sodium Sulfamerazine Activity
78 grams Sodium Sulfamethazine Activity
39 grams Sodium Sulfaquinoxaline Activity

CAUTION: Not for use in humans. Keep out of reach of children. Federal law prohibits the extralabel use of this product in lactating dairy cattle.

Store between 20°C - 25°C (68°F - 77°F) with excursions permitted between 15°C - 40°C (59°F - 104°F).



Restricted Drug (California) - Use only as directed.

Approved by FDA under NADA # 100-094
P08-9001BF Rev. 01-2022

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Huvepharma, Inc. at 1-877-994-4883 or www.huvepharma.us. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

HUVEPHARMA and PoultrySulfa are registered trademarks of Huvepharma EOOD.

CORNER RADIUS

DIRECTIONS

Acute Fowl Cholera - TURKEYS AND CHICKENS: As an aid in the control of acute fowl cholera caused by *Pasteurella multocida* susceptible to sulfamerazine, sulfamethazine and sulfaquinoxaline. Provide medicated water (.04% solution) for 2-3 days. If disease recurs, repeat treatment.

Coccidiosis - TURKEYS: As an aid in the control of coccidiosis caused by *Eimeria meleagridis* and *E. adenoides* susceptible to sulfamerazine, sulfamethazine and sulfaquinoxaline. Provide medicated water (.025% solution) for 2 days, then plain water for 3 days, then medicated water (.025% solution) for 2 days, then plain water for 3 days, then medicated water (.025% solution) for 2 days. Repeat if necessary. DO NOT CHANGE LITTER.

Coccidiosis - CHICKENS: As an aid in the control of coccidiosis caused by *Eimeria tenella* and *E. necatrix* susceptible to sulfamerazine, sulfamethazine and sulfaquinoxaline. Provide medicated water (.04% solution) for 2-3 days, then plain water for 3 days, then medicated water (.025% solution) for 2 days. If bloody droppings appear, repeat at .025% level for 2 more days. DO NOT CHANGE LITTER.

PoultrySulfa®
(sulfamerazine, sulfamethazine and sulfaquinoxaline)

NO COPY